

THE WILSON BULLETIN

A QUARTERLY MAGAZINE OF ORNITHOLOGY

Published by The Wilson Ornithological Society

VOL. 86, No. 2

JUNE, 1974

PAGES 97-196

HEMISPINGUS PARODII, A NEW SPECIES OF TANAGER FROM PERU

JOHN S. WESKE AND JOHN W. TERBORGH

In the course of field work on the ecology and distribution of birds in the eastern Andes of Peru, we encountered a tanager that appears to represent a new species, as described below (and see frontispiece).

Hemispingus parodii sp. nov.

PARODI'S TANAGER

HOLOTYPE.—American Museum of Natural History No. 810,463; Adult male from the Cordillera Vilcabamba, 12°36'S, 73°29'W, elevation 3,480 m, Dept. of Cuzco, Peru; 28 July 1968; collected by John S. Weske and John W. Terborgh; prepared by Weske; original No. 1849.

DIAGNOSIS.—A moderately large *Hemispingus*, nearest *H. calophrys* (Sclater and Salvin, 1876) of the Bolivian Yungas but throat yellow; superciliary stripe yellow and decidedly narrower; crown, lores, and face patch dusky olive; nuchal area with a less pronounced demarcation in color between crown and back; back slightly darker greenish olive; and abdomen more yellowish.

DISTRIBUTION.—At elevations from about 3,200 to 3,500 m on the crest and upper slopes of the northern Cordillera Vilcabamba in the Province of La Convención, Department of Cuzco, Peru. The type locality lies on a high ridge bordering the canyon of the Río Mapitunari, a tributary of the Río Apurímac.

DESCRIPTION OF THE HOLOTYPE.—Back, rump, and upper tail coverts between Citrine and Dark Citrine (capitalized colors are from Ridgway, 1912); crown dusky olive, darker on forehead and lighter posteriorly, the individual feathers being dull blackish with Dark Citrine edgings and basal areas; chin and throat Lemon Chrome; malar area and anterior breast between Lemon Chrome and Primuline Yellow; flanks and sides of breast Citrine; central breast and belly near Lemon Chrome but slightly duller; under tail coverts Sulphine Yellow. A Lemon Chrome superciliary stripe from the base of the bill to the nape, brightest over the eye, the individual feathers being finely tipped with olive; lores and postocular region olive; auriculars and suboculars dull yellowish with paler shaft streaks; postauricular patch Dark Citrine, bordered posteriorly by a dull yellowish streak extending from the throat. Remiges dusky; primaries edged with Sulphine Yellow; secondaries, tertials, and upper wing coverts with margins the color of the back; rectrices with Dark Citrine outer webs and dusky inner webs; tail moderately

graduated, the outermost rectrices being ca. 11 mm shorter than the central (and longest) ones. Soft part colors (in life): maxilla black; mandible medium gray; tarsi and toes pale gray; and iris dark brown.

MEASUREMENTS OF THE HOLOTYPE (mm).—Chord of wing 72.7; tail (from insertion of the two central rectrices to the tip of the longest rectrix) 69.9; tarsus 22.4; bill length (from anterior edge of nostril to tip of maxilla) 7.8; bill depth (at posterior edge of the nostril) 4.9.

ETYMOLOGY.—It is a pleasure to name this species in honor of our friend José Parodi Vargas, who provided us with a base for our expeditions at his Hacienda Luisiana and generously aided us in countless other ways.

REMARKS

Variation in the Type Series.—The eight specimens at hand vary individually in darkness of crown pigmentation but otherwise are quite uniform. One male is blackish on the forecrown and dusky on the occiput, while at the other extreme, another male has a dusky forecrown and a hindcrown which is mainly olive and only slightly darker than the back. On one female, the dusky centers of the crown feathers form fairly distinct streaks, but in the other the crown appears unstreaked and closely matches that of the holotype. Females average smaller than males but otherwise are externally indistinguishable. There is no significant variation in soft part colors.

Habitat.—The habitat of *Hemispingus parodii* is a mixture of elfin forest and tall grassland on the uppermost slopes of the northern Cordillera Vilcabamba. Treeline is ill-defined in this area, but with increasing elevation there is progressively more grassy vegetation and a reduced number of woody plants. Trees are short and gnarled, with a profuse epiphytic covering including lichens, orchids, and bromeliads. Several kinds of bamboo and one species of tree fern are found also. The area is one of cool temperatures and high humidity, for the slopes are customarily in the clouds by day except in early morning. We found *H. parodii* from the summit of the range at 3,520 m down to an elevation of 3,190 m. The general habitat is the same as that of *Gallinago imperialis* and was pictured by Terborgh and Weske (1972).

Breeding Information.—Data on the gonadal condition in our series of *H. parodii* provide a basis for tentatively determining the species' breeding season. Two males taken in early August were in full breeding condition, each having 10 mm testes and a cloacal protuberance (1 and 5 August). The testes of four males collected in mid and late July were considerably enlarged (15 July—6 mm testes; 15 July—9 mm; 21 July—9 × 5 mm; 28 July—left testis 9 × 5 mm and right, 7 × 5 mm). None of these birds had a cloacal protuberance. A female taken 24 July was in breeding condition (ovary and oviduct much enlarged, largest ovum 6 mm) but two August birds were not (2 August—ovary 4 mm, not enlarged; 10 August—ovary 5 mm, largest ovum 1½ mm). These data suggest that the breeding season

for *H. parodi* was just beginning in late July. That breeding had not occurred earlier is suggested by the absence of any juvenal birds or birds with incompletely pneumatized skulls, by the lack of females with a brood patch, and by the relatively unworn plumage of our specimens. Unfortunately we have no information on vocalizations or other behavior.

Ecological Relationships.—A feature of avian distribution on the slopes of the Cordillera Vilcabamba is elevational replacement of ecologically similar species. In accordance with the competitive exclusion principle, evidence points to competition between congeners as a mechanism limiting their respective elevational ranges (Terborgh, 1971). In the northern Cordillera Vilcabamba, *Hemispingus parodii* appears to replace the species *H. atropileus* in the highest montane regions. *H. atropileus*, a slightly larger bird, is common in elfin forest from 2,600 to 2,800 m and was found once at 3,300 m, above the lower elevational limit of *parodii* and in the same netline in which several *parodii* were caught. On the west side of the Apurímac Valley, where *parodii* is not present, *atropileus* ranges to the upper limit of trees. Seemingly, it has been able to expand its elevational range in the absence of competition from its congener. Two much smaller congeneric species, *H. superciliaris* and *H. xanthophthalmus*, have elevational ranges which overlap those of *parodii* and *atropileus* in the Cordillera Vilcabamba, but we feel that they are sufficiently different in size and feeding behavior to minimize competition with the latter two.

All individuals of *H. parodii* that we encountered were trapped in mist-nets. Birds of this species comprised 1.3 percent of our captures in nets from 3,170 to 3,340 m elevation and 0.5 percent of the catch at 3,500 m. Based on capture rates, the commonest species in the area were the tanagers *Iridosornis reinhardti* and *Anisognathus igniventris*, the hummingbird *Metallura tyrianthina*, the flower-piercer *Diglossa lafresnayii*, the finch *Catamenia homochroa*, the furnariid *Schizoeaca fuliginosa*, and the flycatcher *Ochthoeca frontalis*.

Another fairly common species at these elevations is *Basileuterus luteoviridis*. This large warbler bears a striking similarity to *Hemispingus parodii* in color and size. The two species are alike in having citrine upperparts, yellow underparts and superciliaries, and olive facial markings. They are most readily distinguished by crown color—citrine, like the back, in *B. luteoviridis* but dusky olive in *H. parodii*. In addition, *B. luteoviridis* has flesh-colored tarsi and toes, longer rectal bristles, and a flatter bill with a solid black mandible. *B. luteoviridis* is a bird of dense cloud forest understory as well as elfin forest. In the Cordillera Vilcabamba its elevational range is much broader than that of *H. parodii*, extending from 2,100 m to 3,520 m.

TABLE 1
MEASUREMENTS IN MILLIMETERS OF THREE SPECIES OF *HEMISPINGUS*

Measurement	<i>H. parodii</i>				<i>H. calophrys</i>				<i>H. atropileus auricularis</i>			
	N	Mean	S.D.	Range	N	Mean	S.D.	Range	N	Mean	S.D.	Range
MALES												
Wing chord	6	72.0	0.9	70.8-73.0	7	70.0	3.4	65.8-73.9	15	75.6	3.5	70.1-80.4
Tail length	6	68.4	1.4	66.0-69.9	7	66.8	2.0	63.3-69.2	15	72.5	3.0	67.3-79.0
Bill length	6	7.8	0.2	7.5-8.1	7	7.7	0.4	7.2-8.4	14	8.7	0.3	8.0-9.2
Bill depth	6	4.8	0.2	4.5-5.1	5	5.0	0.2	4.8-5.3	14	5.5	0.2	5.0-5.7
FEMALES												
Wing chord	2	69.0	—	67.5-70.5	4	67.4	2.4	64.4-70.1	10	72.9	4.2	65.1-80.5
Tail length	2	66.4	—	65.8-66.9	4	65.6	1.8	63.1-66.9	9	71.4	3.6	65.2-77.9
Bill length	2	8.2	—	8.0-8.3	3	7.6	—	7.0-7.9	10	8.7	0.2	8.5-9.0
Bill depth	2	5.1	—	4.8-5.3	3	4.9	—	4.9-5.0	9	5.4	0.2	5.2-5.7

Symbols: N—sample size, S.D.—standard deviation.

The significance, if any, of the apparent convergence in the two species is not evident.

Systematic Relationships.—In investigating the systematic relationship of *Hemispingus parodii*, we have been led to review the status of the races of *H. atropileus*, which shows considerable resemblance to the new form. Hellmayr (1936) considered *H. atropileus* to consist of three or possibly four subspecies: nominate *atropileus* of eastern Colombia and Ecuador; *chlorigaster*, a western Colombian race perhaps not distinct from *atropileus*; *auricularis*, a somewhat smaller Peruvian form with brighter underparts but like the nominate race in having a narrow, whitish superciliary line; and *calophrys* of Bolivia, a still smaller form with a broad, golden ochraceous superciliary. The race *intermedius*, described by Carriker (1934) from northern Peru, was considered by Zimmer (1947) to be indistinguishable from *auricularis*. It is clear that *H. parodii* and *H. atropileus auricularis* are not conspecific, since their ranges meet and overlap slightly, as was shown above, and there is no indication of intergradation. Since *parodii* and *calophrys* are allopatric, the possibility exists that they are conspecific. Such a taxonomic arrangement would entail recognition of *calophrys* as a species distinct from *H. atropileus*. The habitat, climatic, and elevational preference of *calophrys* is apparently similar to that of *auricularis* and *parodii*. Elevations noted on *calophrys* specimen labels are 3,500 m, 10,000 ft, and 10,700 ft. Bond and Meyer de Schauensee (1942) describe two of its localities with terms like "cold, foggy, and rainy" and "forest and boggy meadows." We have been able to locate only 16 specimens of *calophrys*, all from the Yungas

TABLE 2

COMPARISON OF PLUMAGE COLORATION AND PATTERN IN THREE SPECIES OF *HEMISPINGUS*

Character	<i>H. parodii</i>	<i>H. calophrys</i>	<i>H. atropileatus auricularis</i>
Crown patch			
Width	Wide	Narrow	Wide
Color	Dusky olive	Black	Black
Superciliary stripe			
Width	Narrow	Wide	Narrow
Color	Yellow	Orange-ochraceous	White or buffy-white
Face patch			
Width	Narrow	Narrow	Wide
Color	Dark olive	Blackish to dusky	Black
Throat color	Yellow	Orange-ochraceous	Ochraceous yellow
Color of central abdominal area	Yellow tinged with olive	Yellowish olive	Yellow tinged with olive

of Bolivia in the Departments of La Paz and Cochabamba. We have examined all of these except the holotype, from Tilo Tilo and now in the British Museum, and two individuals, one from Sandillani and one from the La Paz vicinity, now in Munich. No information on the biology of *calophrys* is available in the literature to aid in determining its taxonomic status.

The range of *parodii* and the nearest locality for *calophrys*, not far northeast of La Paz, lie about 750 km apart. The intervening area of Andean slope in southeastern Peru has been visited by such collectors as Chapman, Watkins, and Carriker, but no specimens resembling either of the two forms have been taken. The richness of the black and golden-ochre head coloration varies within the series of *calophrys* specimens at hand, but no morphological trend toward *parodii* in specimens from geographically nearer localities is apparent. Similarly, there is a gap of 570 km between the southernmost locality for *H. atropileus auricularis* (Occobamba Valley, Depto. Cuzco) and the nearest point in the range of *calophrys*. Measurements and characteristics of *parodii*, *calophrys*, and *auricularis* are summarized in Tables 1 and 2. *H. calophrys* seems to differ about equally from *H. parodii* and *H. atropileus auricularis* in plumage coloration and markings. In size, it is similar to the former and somewhat smaller than the latter. Until additional information is available on vocalizations of the various forms and on populations in geographically intermediate areas, if any, it seems best to consider *H.*

parodii and *H. calophrys* each as full species and to recognize *H. atropileus* as a polytypic species (comprising populations to which have been applied the names *chorigaster*, *intermedius*, *auricularis*, as well as *atropileus*), ranging from Colombia to Peru near the city of Cuzco.

SPECIMENS EXAMINED

The museums housing the specimens are cited in parentheses: AMNH, American Museum of Natural History; USNM, U.S. National Museum of Natural History; ANSP, Academy of Natural Sciences of Philadelphia; NS, Naturmuseum Senckenberg; LSU, Louisiana State University Museum of Zoology; FM, Field Museum of Natural History.

Hemispingus parodii.—PERU: Cordillera Vilcabamba (all from within 2 km of the type locality), 6 ♂♂, 2 ♀♀, 1♀ skeleton (AMNH), 1 unsexed alcoholic (USNM).

Hemispingus calophrys.—BOLIVIA: Hichuloma, 1 ♂, 1 ♀ (ANSP); Incachaca, 1 ♂ (ANSP); Cillutincara, 1 unsexed (AMNH); Unduavi, 1 ♂, 1 ♀ (NS); Km 104, Prov. Chapare, 4 ♂♂, 2 ♀♀, 1 unsexed (LSU).

Hemispingus atropileus auricularis.—PERU: Cordillera Vilcabamba, 2 ♂♂, 2 ♀♀ (AMNH); Puncu, Depto. Ayacucho, 1 ♂ (AMNH); Yuracayacu, Depto. Ayacucho, 2 ♀♀ (LSU); Tocopoqueu, Occobamba Valley, 1 ♂ (AMNH), 1 ♂ (USNM); Maraynioc, 2 ♂♂, 1 ♀ (AMNH), 1 ♂ (USNM); Rumicruz, 1 ♂, 2 unsexed (AMNH); Cordillera Carpish, 3 ♂♂, 1 ♀, 1 unsexed (LSU); Carpish Pass, 1 ♀ (AMNH); Carpish Ridge, 1 ♂, 1 ♀ (FM); Bosque Zapatagocha, Depto. Huánuco, 1 ♂ (FM); La Lejia, 1 ♂, 2 ♀♀ (AMNH).

ACKNOWLEDGMENTS

We are grateful to the following curators for loans of specimens and for permission to examine collections in their care: Dean Amadon of the American Museum of Natural History, Emmet R. Blake of the Field Museum of Natural History, Frank B. Gill of the Academy of Natural Sciences of Philadelphia, George H. Lowery, Jr., of the Louisiana State University Museum of Zoology, and Joachim Steinbacher of the Naturmuseum Senckenberg. Information on specimens in the Zoologische Sammlung des Bayerischen Staates was kindly supplied by Gerd Diesselhorst. Eugene Eisenmann and Kenneth C. Parkes critically read the manuscript and made a number of helpful suggestions. Our expeditions to Peru were supported by the Frank M. Chapman Memorial Fund, the National Geographic Society, and the National Science Foundation (Grant No. GB-12378).

LITERATURE CITED

- BOND, J., AND R. MEYER DE SCHAUNSEE. 1942. The birds of Bolivia. Part 1. Proc. Acad. Nat. Sci. Philadelphia. 94:307-391.
- CARRIKER, M. A., JR. 1934. Description of new birds from Peru, with notes on the nomenclature and status of other little-known species. Proc. Acad. Nat. Sci. Philadelphia, 86:317-334.
- HELLMAYR, C. E. 1936. Catalogue of birds of the Americas and adjacent islands. Part 9. Tersinidae-Thraupidae. Field Mus. Nat. Hist., Zool. Ser., 13.

- RIDGWAY, R. 1912. Color standards and color nomenclature. Published by the author, Washington, D.C.
- SCLATER, P. L., AND O. SALVIN. 1876. On new species of Bolivian birds. Proc. Zool. Soc. London, 1876:351-358.
- TERBORGH, J. 1971. Distribution on environmental gradients: theory and a preliminary interpretation of distributional patterns in the avifauna of the Cordillera Vilcabamba, Peru. Ecology, 52:23-40.
- TERBORGH, J., AND J. S. WESKE. 1972. Rediscovery of the Imperial Snipe in Peru. Auk, 89:497-505.
- ZIMMER, J. T. 1947. Studies of Peruvian birds. No. 52. Amer. Mus. Novitates, No. 1367.

BIRD AND MAMMAL LABORATORIES, BUREAU OF SPORT FISHERIES AND WILDLIFE, NATIONAL MUSEUM OF NATURAL HISTORY, WASHINGTON, D.C. 20560, AND DEPARTMENT OF BIOLOGY, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08540. ACCEPTED 28 JANUARY 1974.

NEW LIFE MEMBER



Donald Richard Eckelberry is now a Life Member of the Wilson Ornithological Society. He was born in Sebring, Ohio, and at an early age developed a love of nature and later a facility for painting it. He studied at the Cleveland Institute of Art, and after he left there he pursued a number of jobs before becoming a premier artist and illustrator of birds. He has travelled widely in pursuit of birds, for he prefers to paint species that he knows. Among his best-known feats of illustration are those in Richard Pough's *Audubon Bird Guides* and James Bond's *Guide to the Birds of the West Indies*. Mr. Eckelberry has written several articles on nature and is a member of various organizations, including the A.O.U. He is married and lives in Babylon, Long Island, New York.